# Christopher J. Franklin, D.V.M. Ashley Robinson, D.V.M.



Teresa Petterson, D.V.M. Justin Green, D.V.M.

822 Spring Creek Road · Montrose, CO 81403 Telephone (970) 249-4490 · Fax (970) 240-4580

## POST-OPERATIVE PHYSICAL THERAPY FOR F.H.O. SURGERY

Give medications as directed.

Limit activity for 4 weeks (on leash).

Please contact the hospital if any questions or problems arise.

# Week 1 (at least twice daily)

- 1. Ice the hip region for 5 minutes.
- 2. Gentle massage of thigh and lower leg muscles.
- 3. Flexion and extension of the joints starting at the toes and working up the leg.
- 4. Gentle forward and back movement of the hip region.
- 5. Slow leash walks.
- 6. Recheck appointment 1 week post-op.

#### Week 2

- 1. Continue range of motion exercises, and ice hip as necessary.
- 2. Continue limited activity, increase to 15 minute walks.
- 3. Warm towels can applied to the muscles if there is difficulty stretching.

## Weeks 3 & 4

- 1. Continue controlled exercise, at this point you are trying to regain muscle and strength.
- 2. Repeated short duration exercises are best (e.g. circling, sit/stand, figure-of- eight, etc.)

Due to muscle atrophy, extensive physical therapy is necessary for a successful outcome.

Most patients will permanently have an abnormal gait following this surgery. The purpose of this surgery is to attempt to decrease the severity of pain and increase ambulation.